

PARACETAMOL



commonly used medicine for reducing fever and pain

LIKE ALL MEDICINES, PARACETAMOL CAN CAUSE SIDE EFFECTS

Very rarely, it may cause severe skin reactions, including in children



STAY ALERT

Watch out for these early signs and symptoms

skin reddening







rash



eye irritation



WHAT TO DO?



If any of these signs and symptoms occur, STOP using the medicine immediately and seek medical attention right away.

WHERE TO REPORT?

Please report ANY side effects experienced.

You may report:

- · through your pharmacist or doctor, OR
- directly to NPRA by post or email at fv@npra.gov.my. Scan the QR code to download the reporting form.

Your reports will help the NPRA to take steps to NPRA to take risk.



fv@npra.gov.my

National Pharmaceutical Regulatory Agency Ministry of Health, Malaysia


PARACETAMOL: Reports of Fixed Drug Eruptions

Paracetamol is very rarely associated with fixed drug eruptions (FDEs).

total reports in Malaysia*

41%

in children aged below 12 years



* From Jan 2005 - Mar 2024

PLEASE REMIND PATIENTS (OR THEIR CAREGIVERS):

- ✓ to watch out for these early signs and symptoms of FDE: skin reddening, blisters, rash.
- √ to STOP using the medicine immediately and seek medical attention if these signs and symptoms occur.



REPORT

Please report any adverse events to the NPRA.



National Pharmaceutical Regulatory Agency Ministry of Health, Malaysia



Any queries?

Please send us an email: fv@npra.gov.my